

Growth parameters and mortality rates of Whipfin silver-biddy (*Gerres filamentosus*) and Longtail silver-biddy (*Gerres longirostris*) from creeks of Hara protected area in Jask, Iran

Hashemi S.H.^{*(1)}; Salarpouri A.⁽²⁾; Kiamrani E.⁽³⁾ and Danehkar A.⁽⁴⁾

s.hassanhashemi@yahoo.com

1-Islamic Azad University, Bandar Abbas Branch, P.O.Box: 79145-1311 Bandar Abbas, Iran

2- Persian Gulf and Oman Sea Ecology Research Center, P.O.Box: 1597 Bandar Abbas, Iran

3- Department of Fisheries and Marine Biology, Hormuzgan University, P.O.Box: 3995 Bandar Abbas, Iran

4- Faculty of Natural Resources, Tehran University, P.O.Box: 4111 Karaj, Iran

Received: February 2011

Accepted: January 201

Keywords: Stock assessment, Population dynamic, Gerridae, Oman Sea

Abstract

Population parameters of Whipfin silver-biddy (*Gerres filamentosus*) and Longtail silver-biddy (*Gerres longirostris*) were studied from January to December 2006 in three creeks (Markazi, Khalasi and Gabrik) of Hara protected area in Jask, Oman Sea. All fishes were sampled by boat trawl and beach seine. Growth parameters were determined using the monthly length–frequency parameters. The Mean total length was calculated at 6.2 and 5.4cm for Whipfin silver-biddy and Longtail silver-biddy. The values of L_{∞} and K for Whipfin silver-biddy were 16.8cm and 0.99 yr^{-1} and for Longtail silver-biddy, these were 15cm and 0.77 yr^{-1} , respectively. The value of t_0 was calculated at -0.12 year for both species. The longevity was estimated to be 3 years for Whipfin silver-biddy and 3.8 years for Longtail silver-biddy. The length-weight relationship were determined as $W = 0.0088 L^{3.21}$ and $W = 0.0087 L^{3.18}$ for Whipfin silver-biddy and Longtail silver-biddy, respectively. Results showed the fish were in their juvenile stage which emphasized the supporting roles of creeks for Gerridae as nursery ground.

* Corresponding author